
Species composition and diel variation of fish caught by traditional longline fishing

Panchan, R.^{1*}, Kasamesiri, P.¹, Sutthi, N.¹, Ounjit, W.², Jorjong, S.³ and Sangsila, A.⁴

¹Faculty of Technology, Mahasarakham University, Kantarawichai, Maha Sarakham, Thailand; ²Faculty of Humanities and Social Sciences, Mahasarakham University, Kantarawichai, Maha Sarakham, Thailand; ³Faculty of Natural Resources, Rajamangala University of Technology Isan, Sakon Nakhon Campus, Sakon Nakhon, Thailand; ⁴Division of General Science, Faculty of Education, Buriram Rajabhat University, Buriram 31000, Thailand.

Panchan, R., Kasamesiri, P., Sutthi, N., Ounjit, W., Jorjong, S. and Sangsila, A. (2022). Species composition and diel variation of fish caught by traditional longline fishing. International Journal of Agricultural Technology 18(5):2135-2146.

Abstract The study of fish species composition and diel variation of fish caught by traditional longline fishing was conducted during December 2020 to January 2022. Many sizes of circle hooks ranging from No. 9/0 – No. 18/0 attached with branch lines, main lines and baits (earthworm and small live fish) were used to collect fish samples from 3 stations along the middle Chi River, Maha Sarakham, Thailand. In the current study, a total 414 fish belonging to 9 families with 21 species were found. *Pangasius larnaudii* was the most dominant species in terms of number (15.2%) whereas *Chitala ornate* was the most dominant species in terms of weight (35.6%). The most dominant species in terms of %IRI was *Hemibagrus wyckioides* (16.7%). Total catch per Unit effort of longline fishing ranged from 0.01 – 15.6 kg•day⁻¹ with an average by 2.8 ± 0.31 kg•day⁻¹. Investigation of diel variation of caught fish during December 2021 to January 2022 found that *Hemibagrus wyckioides*, *Hemibagrus filamentus*, *Pangasius larnaudii* and *Mastacembelus armatus* were caught during both day and night. Four and 14 fish species were found during the day and night, respectively. Catch per unit effort during the night with an average of 1,845.5 ± 409.1 g. •12 hours⁻¹ was significantly greater than the catch per unit effort during the day with an average of 264.4 ± 132.7 g. •12 hours⁻¹ (p< 0.05). These results suggested that longline fishing should be operated during the night to gain more catch compared to fishing during the day.

Keywords: Longline fishing, Hook, bait, Species composition, Diel variation, CpUE

Introduction

Fishing is a human activity that is crucial to human life, culture, employment, and economy (Anticamara *et al.*, 2011). Many types of fishing gears have been continually invented and developed associated with fishing area, seasons, size and behavior of target species to increase catch efficiency,

* **Corresponding Author:** Panchan, R.; **Email:** ruamruedee.p@msu.ac.th

reduce bycatch and negative impact of fishing environment (He *et al.*, 2021; Ward and Hindmarsh, 2007). Longline is one of the most used fishing gears worldwide originally developed by Japanese (Watson and Kerstetter, 2006). There are many types of long line according to the target species. These include pelagic longline (drifting longline), bottom longline and (Domingo *et al.*, 2014; He *et al.*, 2021). The main reason of selection this fishing gear that in simply construction, low cost (Bose *et al.*, 2017) and mainly catch larger fishes with high market price (Gilman *et al.*, 2020). Each longline set consists of mainline, branch lines, and baited hooks. The mainline is horizontal line with considerable length depending on fishing area. The branch lines which also called snood or gangions are connected to the mainline with certain interval depending on the size of hook and the target fishes. Baited hooks were attached to the end of each branchline (Hovgard and Lassen, 2000). Longline can be used to fish both day and night times, mostly during the night (Poisson *et al.*, 2010).

Although, longline fishing has been investigated for many years, mostly focused on commercial marine fisheries which are mainly fished for tuna (*Thunnus* spp.) (Domingo *et al.*, 2014; Huang, 2011), swordfish (*Xiphias gladius*) (Amorim *et al.*, 2015), billfishes (*Istiophoridae* spp.) (Gilman *et al.*, 2006; WCPFC, 2019). The investigation of longline fishing within inland fisheries are little known even the longline fishing gear is widely used in many countries like lower Sakarya River, Turkey (Reis and Cerim, 2020), River Tawa, India (Bose *et al.*, 2017).

The Chi River is the longest river in Thailand, approximately 765 km. This river is used for many purposes by surrounding people like agriculture, transportation and irrigation in particular for fishery. Fishing by local fishermen along the river has a long history through the diversity of fishing gears. They have developed and applied their gears with different characteristics and practices related to the season, species, and size of animals. The popular fishing gears were gillnets, lift nets, long line hooks, traps, and surrounding nets (Aengwanich, 1998). Many authors surveyed and reported the fish species diversity caught by many fishing methods, mainly gillnets (Aengwanich *et al.*, 1999; Aengwanich *et al.*, 1998; Leeraputhana, 1997; Nachaiperm *et al.*, 2004; Panchan *et al.*, 2013; Pilasemorn *et al.*, 2006). Investigation of longline fishing in the Chi River is still unknown.

The study aimed to document the long line fishing operation, investigate the catch composition and the dominant fish species of longline fishing and compare catch per unit effort among hook sizes and between the day and the night. The result of this study will provide useful information on the fishery resources to support data related to sustainable management.

Materials and methods

Study sites and fishing gear

The current study was performed between December 2020 and January 2022. All data were collected from 3 sampling sites along a 20 km long section of middle Chi River, Mahasarakham province, the north-eastern region of Thailand (between $16^{\circ} 11' 43''$ to $16^{\circ} 12' 57''$ N and $103^{\circ} 13' 27''$ to $103^{\circ} 19' 22''$ E) (Figure 1).




Figure 1. Sampling sites located at the middle Chi River (S1: Ban Huaychun, S2: Ban Nontoom and S3: Ban BungKha)

The current study was approved the license of the animal welfare No. IACUC-MSU-02/2021 by Mahasarakham University. Local fishermen at each station were assigned to be the data collectors. Before the project started, data collectors were trained on what the aims of the study and the obtained data. The study consisted of 2 parts: 1) interview with the fishermen 2) fishing evaluation.

For the first part, we interviewed the fisherman during fishing operating to gather information about the method and technical characteristics of longline fishing. The second part is to obtain the total catch composition and diel variation of caught fish between day and night. The J-shaped hook with non-offset and 5 different sizes namely 9/0, 12/0, 14/0, 16/0 and 18/0 which were the commonly used by fishermen in the Chi River were used. Two longline sets per hook size (10 sets/ site) were installed for sampling the fish in each study site (Figure 2). The same fishing method and operation were performed at

each sampling sites. Each longline consisted of multifilament mainline of 3.2 mm. attached with monofilament branch lines. Each branch line was connected the baited hook, earthworm, or small alive fish as bait. The fishermen were free decision when to fish on each appointment fishing date. Total 59 days were the fishing days of the current study. On each fishing days, the fisherman at each sampling site set the long lines during 3-6 pm. The gears were then left to fish for about 12 hours. In the morning of the next day at 6-9 am, they came back to collect the fish from each set of longline. For the investigation of diel variation in catch composition, we performed the experiment at site 1 between December 2021 to January 2022. The gears were set for 24 hours in fishing area and fish samples were collected every 12 hours. The fish samples of both experiments were taxonomically identified into species (Rainboth, 1996), counted, measured the total length (nearest 0.1 cm.) and weighted (nearest 0.1g.) in each hook size and each fishing period.




Figure 2. J-shaped hooks in different sizes ranging from 9/0 - 18/0 are commonly used in the Chi River

Data analysis

From the data the percentage of catch composition in term of family was analysed. Percentage weight, number and occurred frequency of each fish species in the total catch were used to calculate the index of relative importance (%IRI) (Pinkas *et al.*, 1971) as follow equation:

$$\%IRI = \left(\frac{(\%W_i + N_i) \times \%F_i}{\sum_{i=1}^n (\%W_i + \%N_i) \times \%F_i} \right) \times 100$$

Where $\%W_i$ and $\%N_i$ are the percentage weight and number of the i^{th} species in the total catch and $\%F_i$ is the percentage frequency of occurrence of each species in the total number of species.

The catch per unit effort (CpUE) was expressed as kg.day^{-1} by each hook and kg.12 hrs^{-1} by each fishing period for 12 hours. Kruskal-Wallis chi-square test as non-parametric statistic was used to test the equal of median of CpUE among group of hook sizes and the fishing periods. Tukey's HSD was posteriori test when the difference among group of hook size was found. Wilcoxon-Mann-Withey U- test, a non- parametric statistic was used to test the different of median between the fishing period.

Results

Fishing operation

Firstly, fisherman installed the main line across the river and holded the line with either stone or tree at the riverbanks (Figure 3). The length of main line depended on the river width approximately 50-70 m. Secondly, each branch lines with length 1-2 m. with the baited hook were connected to the main line with certain interval between line approximately 2 - 4 m. Baits are usually the small alive fish and the earthworm (Figure 3). There were 10-15 hooks per long line set. Stones connecting the other lines were connected to the main line to set the depth of longline in the water during fishing. The depth of longline was relative the distance between the stone and mainline. The shorter distance was deeper in the water than the longer distance. The fisherman usually set the long lines in dusk at 3-6 pm. The gears were then left to fish for about 12 hours. They came back to collect the fish in the morning of the next day at 6-9 am. For the collecting process, each hook was checked without retrieving the gear from the water. The new bait was re-hooked when the lost bait was found. For collecting the caught fish, mostly the fish were caught by swallowing the baited hook into their mouth. Fishermen carefully removed the fish from the hook without cut off the branchline if the small fish was caught. However, the branchline with hook was probably cut off from the mainline to collect the larger fish on boat if they were unable easily remove the hook. Then, substitution of the new branch line with baited hook was installed. The total time spending of collecting fish was approximately 2-3 hours. The fishermen, moreover, reported that the target species of long line fishing were the larger fishes like *P. larnaudii*, *H. wyckioides* and *C. ornate* which were the economical important species.

Catch composition and the dominant fish species of longline fishing

In total 59 fishing date during December 2020 to January 2022, 414 fish belonging 9 families 17 genera and 21 species were found (Table 1).

Cyprinidae was the most frequent occurrence in terms of family, which comprised 6 fish species (28.6%), followed by Bagridae (19.0%) and Siluridae (14.3%), respectively (Figure 4). *P. larnaudii*, *C. ornata* and *H. wyckioides* were the most dominant species in terms of number at 15.2%, 14.0% and 11.1%, respectively (Table 1). *C. ornata*, *H. wyckioides* and *W. Attu* were the most dominant species in terms of weight at 35.6%, 29.8% and 9.6%, respectively (Table 1). According to percentage index of relative importance (%IRI) *H. wyckioides* was the dominant species caught by longline fishing at 16.7% followed by *P. larnaudii*, *H. filamentus*, *C. ornata* were equally %IRI by 12.8% (Figure 4).

Consideration catch composition by the fishing period during December 2021 to January 2022 (Table 2), *H. wyckioides*, *H. filamentus*, *P. larnaudii* and *M. armatus* were caught during both day and night. Four and 14 fish species were found during the day and night, respectively.




Figure 3. Schematic represents the operation of longline fishing between the riverbanks in the Chi River with using small alive fish and earthworm as bait

Catch per unit effort (CpUE)

The investigation of CpUE among hook size throughout the study period ranged from 0.01 and 15.6 kg• day⁻¹ with an average by 2.8 ± 0.31 kg •day⁻¹ (Table 3). The largest hook size No. 9/0 revealed the greater CpUE than the others hook ($p < 0.05$) (Table 3 and Figure 5a). Consideration CpUE in term of different fishing period, catch per unit effort during the night with an average of $1,845.5 \pm 409.1$ g•12 hours⁻¹ was significantly greater than the catch per unit effort during the day with an average of 264.4 ± 132.7 g. •12 hours⁻¹ ($p < 0.05$) (Table 3 and Figure 5b).

Table 1. Species composition by different hook size and percentage index of relative importance (%IRI) of longline fishing in the Chi River. Bold letters in species column refer to family

Species	Frequency					%	Weight (g)					%	%IRI
	9/0	12/0	14/0	16/0	18/0		9/0	12/0	14/0	16/0	18/0		
Notopteridae													
<i>Chitala ornata</i>	15	12	8	11	-	11.1	41.9	26.6	10.80	3.50	-	35.6	12.8
<i>Notopterus notopterus</i>	1	-	1	-	-	0.5	0.13	-	0.1	-	-	0.1	1.3
Cyprinidae													
<i>Cyclocheilichthys enoplos</i>	8	2	-	9	-	4.6	12.0	2.00	2.95	-	-	7.3	5.1
<i>Puntioplites proctozysron</i>	-	-	2	-	1	0.7	-	-	0.27	-	0.01	0.1	1.9
<i>Barbonymus gonionotus</i>	1	-	3	-	-	1.0	0.02	-	0.59	-	-	0.3	2.6
<i>Probarbus jullieni</i>	-	-	1	-	-	0.2	-	-	1.88	-	-	0.8	0.6
<i>Barbonymus altus</i>	-	-	-	-	3	0.7	-	-	-	-	0.10	0.04	0.6
<i>Labeo chrysophekadion</i>	-	1	-	-	-	0.2	-	1.10	-	-	-	0.5	0.6
Bagridae													
<i>Hemibagrus wyckioides</i>	19	4	32	-	3	14.0	47.80	8.80	7.47	-	5.3	29.8	16.7
<i>Hemibagrus filamentus</i>	2	13	21	-	14	12.1	0.22	2.99	1.65	-	1.56	2.8	12.8
<i>Mystus nigriceps</i>	-	-	2	39	-	9.9	-	-	0.27	5.41	-	2.4	4.5
<i>Mystus bocourti</i>	-	2	-	-	-	0.5	-	0.08	-	-	-	0.03	0.6
Siluridae													
<i>Wallago Attu</i>	10	1	6			4.1	20.8	1.20	0.46	-	-	9.6	5.1
<i>Ompok bimaculatus</i>	1	-	13	-	1	3.6	0.10	-	0.69	0.06	-	0.4	3.2
<i>Micronema micronema</i>	5	7	10	8	3	8.0	1.20	2.03	1.35	1.30	0.58	2.8	10.9
Pangasiidae													
<i>Pangasius larnaudii</i>	13	13	6	9	2	15.2	2.56	4.27	2.45	0.40	0.65	4.4	12.8
<i>Pangasius macronema</i>	-	10	-	-	-	2.4	-	0.13	-	-	-	0.1	0.6
Clariidae													
<i>Clarias batrachus</i>	-	-	1	-	-	0.2	-	-	0.74	-	-	0.32	
Mastacembelidae													
<i>Mastacembelus armatus</i>	-	1	36	-	1	9.2	-	0.11	2.77	-	0.01	1.3	3.8
Eleotridae													
<i>Oxyeleotris marmorata</i>	1	-	3	-	1	1.2	0.20	-	0.21	-	1.00	0.6	1.9
Chaniuidae													
<i>Channa striata</i>	-	2	-	-	-	0.5	-	2.00	-	-	-	0.9	0.6
Total	76	88	145	76	29		126.9	51.3	31.7	13.6	9.3		





Figure 4. Percentage of fish species in each family caught by longline fishing

Table 2. Species composition caught by longline fishing between day and night in the Chi River. Bold letters in species column refer to family

Species	Frequency		%		Weight (g)		%	
	Day	Night	Day	Night	Day	Night	Day	Night
Cyprinidae								
<i>Puntioplites proctozysron</i>	2	0	1.64	0	268	-	1.59	-
<i>Barbonymus gonionotus</i>	3	0	2.46	0	589	-	3.49	-
<i>Probarbus jullieni</i>	1	0	0.82	0	1,880	-	11.14	-
<i>Barbonymus altus</i>	3	0	2.46	0	96	-	0.57	-
Bagridae								
<i>Hemibagrus wyckioides</i>	17	7	13.93	5.74	4,851	323	28.74	1.91
<i>Hemibagrus filamentus</i>	11	5	9.02	4.10	401	165	2.38	0.98
<i>Mystus nigriceps</i>	1	0	0.82	0	67	-	0.40	-
Siluridae								
<i>Wallago Attu</i>	6	0	4.92	0	456	-	2.70	-
<i>Ompok bimaculatus</i>	14	0	11.48	0	746	-	4.42	-
<i>Micronema bleekeri</i>	5	0	4.10	0	530	-	3.14	-
Pangasiidae								
<i>Pangasius larnaudii</i>	5	1	4.10	0.82	2,213	505	13.11	2.99
Clariidae								
<i>Clarias batrachus</i>	1	0	0.82	0	740	-	4.38	-
Mastacembelidae								
<i>Mastacembelus armatus</i>	23	14	18.85	11.48	1,719	1,122	10.18	6.65
Eleotridae								
<i>Oxyeleotris marmorata</i>	3	0	2.46	0	208	-	1.23	-
Total	14	4			14,764	2,115		

Table 3. Catch per Unit effort of longline fishing in the Chi River

Hook size	CpUE (kg•day ⁻¹)		Fishing period	CpUE (g. •12 hours ⁻¹)	
	Min - Max	Average±SE		Min -Max	Average±SE
No. 9/0	0.1-15.6	4.1 ± 0.6	Day	0 – 106	264.4 ± 132.7
No. 12/0	0.2 - 11.7	2.6 ± 0.7	Night	321 – 4000	1.845.5 ± 409.1
No. 14/0	0.3 - 4.9	1.7 ± 0.3	Total	0 – 4000	264.4 ± 132.7
No.16/0	0.8 - 6.9	2.7 ± 1.1			
No.18/0	0.01 - 3.5	1.0 ± 0.4			
Total	0.01 – 15.6	2.8 ± 0.31			

**Figure 5.** Summary of statistical testing of CpUE at different hook size (a) and the fishing period (b). The same letter above the boxes indicate that the values are not statistically different ($\alpha = 0.05$)

Discussion

The results revealed the basic information regarding the catch composition and catch per unit effort of the long line fishing in the Chi River, which have not been investigated ever. The interview about fishing operation,

hook size 9/0, 12/0, 14/0 16/0 and 18/0 with small alive fish and earthworm as bait were mostly used to fish by fishermen. The most important factors effecting long line efficiency were hook size and bait type. The hook size and bait type would be selected for fishing depending on the needed target species (Løkkeborg and Bjordal, 1992). In Chi River, the fishermen reported that the target species of long line fishing were the larger fishes like *P. larnaudii*, *H. wyckioides* and *C. ornate* which were the economical important species.

The results of fishing observation were found 414 fish belonging 9 families 17 genera and 21 species which differ to the previous study (Leeraputhana, 1997; Aengwanich *et al.*, 1998; Nachaiperm *et al.*, 2004; Pilasemorn *et al.*, 2006; Panchan *et al.*, 2013). This might be due to the different fishing method and location to sampling the fishes. However, family Cyprinidae was the pre-dominant group similar to the previous studies. Aengwanich *et al.*, 1999; Nachaiperm *et al.*, 2004; Jutagate, 2009 reported that fish in the Cyprinidae family are the predominant fish species found in the Chi River and are commonly found within Thai riverine habitats.

Longline fishing gear is classified as highly selectivity like the selectivity of gillnets. Both gears were catch specific by species and size. The selectivity of species specific depends on either the fishing strategies or fishin gears including the fish distribution, both horizontal and vetical distributions (Løkkeborg and Bjordal, 1992). The study, the long lines were installed in the mid- water depth. The result revealed that *H. wyckioides*, *P. larnaudii*, *H. filamentus* and *C. ornate* were the dominant species caught by longline fishing. These species live between mid-water and bottom area. The result differed to gillnet fishing, mostly the pelagic fishes were caught (Nachaiperm *et al.*, 2004; Pilasemorn *et al.*, 2006; Panchan *et al.*, 2013).

Catch per unit effort (CpUE) is one of the important indicators on the status of fish abundance related to usable stocks and fishing methods (Bishop, 2006). In the study, CpUE among hook sizes period ranged from 0.01 – 15.6 with an average by $2.8 \pm 0.31 \text{ kg} \cdot \text{day}^{-1}$. Differently, the investigation of traditional longline set used in the lower Sakarya River revealed that the average daily catch was between 5- 10 kg.(Reis and Cerim, 2020). Moreover, the result differed from the previous studies in the Chi River by gillnet like $2.1 \pm 0.4 \text{ kg} \cdot \text{fisherman}^{-1} \cdot \text{day}^{-1}$ (Panchan *et al.*, 2013), $147.80 \text{ g} \cdot 100\text{m}^2 \cdot \text{hr}^{-1}$ (Pilasemorn *et al.*, 2006) and $6.80 \text{ g} \cdot 180\text{m}^2 \cdot \text{hr}^{-1}$ (Nachaiperm *et al.*, 2004). Generally, fishes were caught by swallowing the baited hook in the mouth or penetrating their flesh. The hook characteristic and the biological charactertistic of fish affected the catch performance (Løkkeborg and Bjordal, 1992). Structure of large hook was stronger and gab wider and consequently more effective catch than smaller hook. Likewise, the highest catch per unit effort

was found at hook size 9/0, ranging from 0.11 - 15.6 kg•day⁻¹ (an average 4.1±0.6 kg•day⁻¹).

Catch per unit effort (CpUE) of long line fishing during the night with an average of 1,845.5 ± 409.1 g•12 hours⁻¹ was significantly greater than the catch per unit effort during the day (an average of 264.4 ± 132.7 g •12 hours⁻¹). There are many factors affecting the performance of longline fishing, whether external or internal factors of fish like hunger state, food habit and feed abundance (Løkkeborg *et al.*, 2014). Moreover light level as the external factor affects visibility of fishing gear and bait of the fish (Stoner, 2003), resulting in low CpUE during the day. Fourteen fishes were found in both fishing periods, however, might be due to hunger state of fish.

Acknowledgements

This research project was financially supported by Mahasarakham University 2021, grant No. 6408040/2564. We would like to thank the staff of Department of Agriculture Technology (Fisheries), Faculty of Technology, Mahasarakham University and Mr. Jirapong Ponsen who substantially contributed to the field work. We are grateful to the local fishermen from the Chi River, Mahasarakham province for their kind help during the field trips.

References

- Amorim, S., Santos, M. N., Coelho, R. and Fernandez-Carvalho, J. (2015). Effects of 17/0 circle hooks and bait on fish catches in a Southern Atlantic swordfish longline fishery. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 25:518-533.
- Aengwanich, W. (1998). Fish culture of the Chi River Basin community. The Thailand Research Fund (TRF). 316 pp.
- Aengwanich, W., Chinrasri, T. and Chinrasi, O. (1999). Study on diversity of family Cyprinidae of Chi River in Mahasarakham province, Thailand . *KKU Science Journal*, 27:43-50.
- Aengwanich, W., Chinrasri, T., Chinrasi, O. and Kongbuntad, W. (1998) Ichthyofauna and taxonomy of the Chi River's fish in Mahasarakham province. Mahasarakham.
- Anticamara, J. A., Watson, R., Gelchu, A. and Pauly, D. (2011). Global fishing effort (1950-2010): Trends, gaps, and implications. *Fisheries Research*, 107:131-136.
- Bishop, J. (2006). Standardizing fishery-dependent catch and effort data in complex fisheries with technology change. *Reviews in Fish Biology and Fisheries*, 16:21-38.
- Bose, A. K., Gupta, S., Suresh, V. R. and Das, A. K. (2017). Indigenous fishing (Char Kanta): A traditional fisheries in River Tawa , Madhya Pradesh , India. *International Journal of Fisheries and Aquatic Studies*, 5:265-268.
- Domingo, A., Rodrigo, F., Philip, M. and Jiménez, S. (2014). Longline. In *ICCAT Manual*, pp.1-48. <https://doi.org/10.13140/2.1.1615.8088>
- Gilman, E., Chaloupka, M., Bach, P., Fennell, H., Hall, M., Musyl, M. and Song, L. (2020). Effect of pelagic longline bait type on species selectivity: a global synthesis of evidence. *Reviews in Fish Biology and Fisheries*, 30:535-551.
- Gilman, E., Zollett, E., Beverly, S., Nakano, H., Davis, K., Shiode, D. and Kinan, I. (2006).

- Reducing sea turtle by-catch in pelagic longline fisheries. *Fish and Fisheries*, 7:2-23.
- He, P., Chopin, F., Suuronen, P., Ferro, S. T. R. and Lansley, J. (2021). Classification and illustrated definition of fishing gears. In *Classification and illustrated definition of fishing gears*. Rome: FAO Fisheries and Aquaculture Technical Paper No. 672. <https://doi.org/10.4060/cb4966en>
- Hovgard, H. and Lassen, H. (2000). Manual on estimation of selectivity for gillnet and longline gears in abundance surveys. In FAO. Fisheries Technical Paper. Retrieved from <http://www.fao.org/3/x7788e/X7788E00.htm#TOC>
- Huang, H. W. (2011). Bycatch of high sea longline fisheries and measures taken by Taiwan: Actions and challenges. *Marine Policy*, 35:712-720.
- Jutagate, T. (2009). Reservoir fisheries in Thailand . In: de Silva SS, Amarasinghe US, editors. *Reservoir Fisheries in Asia and Pacific*. Bangkok: Network of Aquaculture Centres in Asia-Pacific (NACA). p.96-113.
- Leeraputhana, W. (1997). *Fish and fisheries in Chi River* . Bangkok.
- Løkkeborg, S. and Bjordal, Å. (1992) Species and size selectivity in longline fishing: a review . *Fisheries Research*, 13:311-322.
- Løkkeborg, S., Siikavuopio, S. I., Humborstad, O. B., Utne-Palm, A. C. and Ferter, K. (2014). Towards more efficient longline fisheries: fish feeding behaviour, bait characteristics and development of alternative baits . *Reviews in Fish Biology and Fisheries*, 24:985-1003.
- Nachaiperm, J., Neungsith, C. and Chernchom, C. (2004). Structure and distribution of fish community in Chi River . Bangkok.
- Panchan, R., Jutagate, T. and Wigraiboon, S. (2013). Fish species composition caught by gillnets: Case study from Chi River, Mahasarakham province, Thailand (in Thai) . *Maejo International Journal of Science and Technology*, 7:43-51.
- Pilasemorn, C., Singsri, S., Phetcharat, C. and Chaengkij, M. (2006). Structure and distribution of fish community in the Chi River during 2003-2005 . Bangkok.
- Pinkas, L., Oliphant, M. S. and Iverson I. L. K. (1971). Food Habits of Albacore, Blue-fin Tuna, and Bonito in California Waters . *Calif Dep Fish Game's Fish Bull*, 152:5-10.
- Poisson, F., Gaertner, J. C., Taquet, M., Durbec, J. P. and Bigelow, K. (2010). Effects of lunar cycle and fishing operations on longline-caught pelagic fish: Fishing performance, capture time, and survival of fish. *Fishery Bulletin*, 108:268-281.
- Rainboth, W. J. (1996). *Fishes of Cambodian Mekong* . FAO. Rome.
- Reis Í, and Cerim H (2020) Method and Technical Characteristics of Traditional River Longline from Lower Sakarya River Fishery, Turkey . *Journal of Limnology and Freshwater Fisheries Research*, 6:164-168.
- Stoner A. W. (2003). Hunger and light level alter response to bait by Pacific halibut: Laboratory analysis of detection, location and attack . *Journal of Fish Biology*, 62:1176-1193.
- Ward, P. and Hindmarsh, S. (2007). An overview of historical changes in the fishing gear and practices of pelagic longliners, with particular reference to Japan's Pacific fleet. *Reviews in Fish Biology and Fisheries*, 17:501-516.
- Watson, J. W. and Kerstetter, D. W. (2006). Pelagic longline fishing gear: A brief history and review of research efforts to improve selectivity. *Marine Technology Society Journal*, 40:6-11.
- WCPFC. (2019). *Western and Central Pacific Fisheries Commission Tuna Fishery Yearbook 2018*.

(Received: 3 August 2022, accepted: 3 September 2022)